



IT'S THEIR TIME TO GROW AND LEARN. TAKE THEIR DAY BACK FROM SEIZURES

Discover how your child may experience fewer partial-onset (focal) seizures by taking BRIVIACT alone or adding it to their current treatment.

INDICATION

BRIVIACT® (brivaracetam) CV is a prescription medicine used to treat partial-onset seizures in people 1 month of age and older.

IMPORTANT SAFETY INFORMATION

BRIVIACT may cause serious side effects including suicidal thoughts or actions; drowsiness; tiredness; dizziness; balance and coordination problems; mood and behavior changes including irritability, anxiety, and psychotic symptoms; and allergic reactions to ingredients in BRIVIACT.

Please see Important Safety Information on pages 12-14 of this brochure and review the [Medication Guide](#) on [BRIVIACT.com](#).

Save on BRIVIACT. See inside for details.

BRIVIACT® 
(brivaracetam) 

MANAGING PARTIAL-ONSET SEIZURES IN YOUR CHILD AT EVERY STAGE OF DEVELOPMENT

A seizure can occur at any age—and when you least expect it.

New cases of partial-onset seizures are common in the first year of life but are unpredictable and can strike at any age. As a parent, you want to be prepared with the right treatment and information.

Whether your child is an infant, toddler, or teen, and whether he or she has been recently diagnosed or struggling for some time, you should know there is hope. You may be able to help reduce the number of partial-onset seizures your child is having.

Use this brochure to help you better understand BRIVIACT® (brivaracetam) CV. BRIVIACT may not be for everyone. Ask your child's healthcare provider if BRIVIACT is right for him or her.

BRIVIACT is a prescription medicine used to treat partial-onset seizures in people 1 month of age and older.

It is not known if BRIVIACT is safe and effective in children younger than 1 month of age.



Did you know?
YOU CAN SAVE ON BRIVIACT.

See page 16 of this brochure
for details.



IMPORTANT SAFETY INFORMATION

Suicidal thoughts or actions: Like other antiepileptic drugs, BRIVIACT may cause suicidal thoughts or actions in a very small number of people, about 1 in 500 people taking it. Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you: thoughts about suicide or dying; attempts to commit suicide; new or worse depression; new or worse anxiety; feeling agitated or restless; panic attacks; trouble sleeping (insomnia); new or worse irritability; acting aggressive, feeling angry, or being violent; acting on dangerous impulses; an extreme increase in activity and talking (mania); other unusual changes in behavior or mood. Before taking BRIVIACT, tell your healthcare provider if you have or had depression, mood problems, or suicidal thoughts or behavior.

Please see Important Safety Information on pages 12-14 of this brochure and review the [Medication Guide](#) on [BRIVIACT.com](#).

What's inside?

Page

UNDERSTANDING EPILEPSY

Learn more about partial-onset seizures and treatment.

04

HOW BRIVIACT CAN HELP

Discover how BRIVIACT may help your child achieve fewer partial-onset seizures.

06

TAKING BRIVIACT

Learn about starting treatment with BRIVIACT.

08

MEDISAFE APP

A resource that can provide medication reminders.

10

IMPORTANT SAFETY INFORMATION

Learn about the possible side effects of BRIVIACT.

12

HELPFUL RESOURCES

Information and tools to help you as a caregiver.

15

SAVE ON BRIVIACT

Use the Patient Savings Card to save on BRIVIACT prescriptions.

16

HAVE MORE QUESTIONS?

If you have any questions or want more information, please contact us.

17

UNDERSTANDING EPILEPSY

As a parent, you know **the impact that partial-onset seizures can have on your child's day**. Understanding epilepsy is an important part of caring for your child so they have the chance to learn, play, and grow.

What is epilepsy?

- Epilepsy is a condition of the brain causing **repeated and unpredictable seizures**
- A seizure is caused by a **disturbance** in the **normal electrical activity of the brain**
- Someone may be diagnosed with epilepsy if they experience **2 or more of these seizures** separated by **at least 24 hours**

Partial-onset seizures are the most common type of seizure in children.

About two-thirds of seizures in children are partial-onset seizures, also known as focal seizures.

Partial-onset seizures:

- Can be serious and can affect your child's ability to respond to what's happening around them
- May be hard to recognize in your child

Go to **[BRIVIACT.com](https://www.briviac.com)** to learn more about the different types of partial-onset seizures and how BRIVIACT may help.

IMPORTANT SAFETY INFORMATION

Nervous system problems: Drowsiness, tiredness, and dizziness are common with BRIVIACT, but can be severe. BRIVIACT can also cause problems with balance and coordination. Do not drive or operate machinery until you know how BRIVIACT affects you.

Please see Important Safety Information on pages 12-14 of this brochure and review the **[Medication Guide](#)** on **[BRIVIACT.com](https://www.briviac.com)**.



Around 3.4 million people in the United States are living with epilepsy.

About **470,000** of them are children.

HOW BRIVIACT CAN HELP

How can BRIVIACT help my child?

BRIVIACT is a prescription medicine that can be used to treat partial-onset seizures in patients 1 month of age and older.

BRIVIACT can be **taken alone or may be added to an existing seizure medicine** to help control partial-onset seizures. BRIVIACT may help reduce the number of partial-onset seizures your child is having.

YOU CAN SAVE ON BRIVIACT.*

Check page 16 of this brochure for details on the BRIVIACT Patient Savings Card.

*Subject to eligibility. Terms and Conditions apply.



IMPORTANT SAFETY INFORMATION

Mental (psychiatric) symptoms: BRIVIACT can cause mood and behavior changes such as aggression, agitation, anger, anxiety, apathy, mood swings, depression, hostility, and irritability. Irritability and anxiety are common with BRIVIACT, and can be severe. People who take BRIVIACT can also get psychotic symptoms such as hallucinations (seeing or hearing things that are really not there), delusions (false or strange thoughts or beliefs), and unusual behavior.

Please see Important Safety Information on pages 12-14 of this brochure and review the [Medication Guide](#) on BRIVIACT.com.



TAKING BRIVIACT

Your child's doctor will prescribe an appropriate dose and formulation of BRIVIACT based on your child's needs. However, it can be challenging to get some kids to take their medicine. That's why BRIVIACT has 2 options for oral formulations: tablets and a raspberry-flavored liquid.



(Not actual size)

BRIVIACT TABLETS

BRIVIACT tablets come in multiple doses and should be taken whole. Do not crush or chew tablets before swallowing.



Raspberry
flavor

BRIVIACT ORAL SOLUTION

If your child has trouble taking pills, he or she can take BRIVIACT as a liquid.



The hope is that a child never has to go to the hospital. But if they do and they can't swallow the oral solution or tablets, a **healthcare professional can give BRIVIACT through the vein (intravenous).**

IMPORTANT SAFETY INFORMATION

Allergic reactions: Do not take BRIVIACT if you are allergic to brivaracetam or any of the ingredients in BRIVIACT. See the Patient Medication Guide included for a list of ingredients in BRIVIACT.

Abuse and dependence: BRIVIACT is a federally controlled substance (CV) because it can be abused or lead to dependence. Tell your healthcare provider if you have abused or been dependent on prescription medicines, street drugs, or alcohol in the past. Keep BRIVIACT in a safe place to prevent misuse and abuse. Selling or giving away BRIVIACT may harm others and is against the law.

Other medical conditions: Before taking BRIVIACT, tell your healthcare provider about all of your medical conditions, including if you have liver problems.

Please see Important Safety Information on pages 12-14 of this brochure and review the [Medication Guide on BRIVIACT.com](#).



Remember this information when giving your child BRIVIACT:

- **Never stop taking BRIVIACT without first talking to a healthcare provider**
 - **Stopping BRIVIACT suddenly can cause serious problems.** Stopping a seizure medicine suddenly can cause seizures that will not stop (status epilepticus)
 - **If your doctor decides to stop your child's BRIVIACT treatment,** he or she will give you instructions on how to slowly stop taking BRIVIACT
- If your healthcare provider has prescribed BRIVIACT oral solution, be sure to **ask your pharmacist for a medicine dropper or medicine cup** to help you measure the correct amount of BRIVIACT oral solution. Do not use a household teaspoon or tablespoon. **Ask your pharmacist for instructions** on how to use the measuring device the right way
- Your child should take BRIVIACT **exactly as his or her doctor has instructed**
- Your child's doctor will tell you **how much** BRIVIACT to take and **when** to take it
- If your child takes too much BRIVIACT, call your local Poison Control Center immediately
- Your child can take BRIVIACT **with or without food**
- BRIVIACT can be stored at **room temperature**

Have more questions? Remember to ask your child's doctor or healthcare team.



NEED HELP MANAGING YOUR CHILD'S MEDICATION? TRY MEDISAFE®

Medisafe is a free app you can use to help your child take their medication exactly as prescribed.

Stay on track and in control with features like:



Personalized medication, refill, and appointment reminders




Helpful resources and more

Two ways to get Medisafe

1. **Download** the Medisafe app at [MedisafeApp.com](https://www.MedisafeApp.com) on your mobile phone



2. **Scan** the QR code below using your phone camera



Record your child's seizures so you can share the information with their doctor at their next appointment.

Go to BRIVIACT.com/tools-resources

Once Medisafe is on your mobile phone, simply:

TAP
"Add your med"

TYPE
"BRIVIACT"

ENROLL
as a caregiver

IMPORTANT SAFETY INFORMATION

Most common side effects: In adult clinical trials, the most common side effects of BRIVIACT include sleepiness, dizziness, feeling tired, and nausea and vomiting. Side effects of BRIVIACT in children 1 month to less than 16 years of age are similar to those seen in adults.

Please see Important Safety Information on pages 12-14 of this brochure and review the [Medication Guide](https://www.BRIVIACT.com) on BRIVIACT.com.

IMPORTANT SAFETY INFORMATION



Be educated about the Important Safety Information and talk to your child's doctor about concerns you may have or should anything out of the ordinary take place.

IMPORTANT SAFETY INFORMATION

What is BRIVIACT?

BRIVIACT® (brivaracetam) CV is a prescription medicine used to treat partial-onset seizures in people 1 month of age and older. It is not known if BRIVIACT is safe and effective in children younger than 1 month of age.

What is the most important information I should know about BRIVIACT?

BRIVIACT is a federally controlled substance (CV) because it can be abused or lead to dependence. Keep BRIVIACT in a safe place to prevent misuse and abuse. Selling or giving away BRIVIACT may harm others and is against the law.

Like other antiepileptic drugs, BRIVIACT may cause suicidal thoughts or actions in a very small number of people, about 1 in 500 people taking it.

Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- new or worse depression
- feeling agitated or restless
- trouble sleeping (insomnia)
- acting aggressive, feeling angry, or being violent
- an extreme increase in activity and talking (mania)
- attempts to commit suicide
- new or worse anxiety
- panic attacks
- new or worse irritability
- acting on dangerous impulses
- other unusual changes in behavior or mood

Do not stop BRIVIACT without first talking to a healthcare provider.

- Stopping BRIVIACT suddenly can cause serious problems.
- Stopping a seizure medicine suddenly can cause seizures that will not stop (status epilepticus).

Who should not take BRIVIACT?

Do not take BRIVIACT if you are allergic to brivaracetam or any of the ingredients in BRIVIACT.

What should I tell my healthcare provider before starting BRIVIACT?

Before taking BRIVIACT, tell your healthcare provider about all of your medical conditions, including if you:

- have or had depression, mood problems, or suicidal thoughts or behavior.
- have liver problems.
- have abused or been dependent on prescription medicines, street drugs, or alcohol.
- are pregnant or plan to become pregnant. It is not known if BRIVIACT will harm your unborn baby.
- are breastfeeding or plan to breastfeed. BRIVIACT passes into your breast milk.

What should I avoid while taking BRIVIACT?

Do not drive or operate machinery until you know how BRIVIACT affects you. BRIVIACT may cause drowsiness, tiredness, dizziness, and problems with your balance and coordination.

What are the possible side effects of BRIVIACT?

BRIVIACT may cause serious side effects, including:

- See **“What is the most important information I should know about BRIVIACT?”**
- **Nervous system problems.** Drowsiness, tiredness, and dizziness are common with BRIVIACT, but can be severe. See **“What should I avoid while taking BRIVIACT?”** BRIVIACT can also cause problems with balance and coordination.
- **Mental (psychiatric) symptoms.** BRIVIACT can cause mood and behavior changes such as aggression, agitation, anger, anxiety, apathy, mood swings, depression, hostility, and irritability. Irritability and anxiety are common with BRIVIACT, and can be severe. People who take BRIVIACT can also get psychotic symptoms such as hallucinations (seeing or hearing things that are really not there), delusions (false or strange thoughts or beliefs), and unusual behavior.

IMPORTANT SAFETY INFORMATION (CONT.)

The most common side effects of BRIVIACT in adults include:

- sleepiness
- dizziness
- feeling tired
- nausea and vomiting

Side effects of BRIVIACT in children 1 month to less than 16 years of age are similar to those seen in adults.

These are not all the possible side effects of BRIVIACT. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to UCB, Inc. at UCBCares (1-844-599-CARE [2273]).

HELPFUL RESOURCES



We're here with the information and tools you need as a caregiver to help you and your child learn about and manage partial-onset seizures. Take a few minutes to get to know the resources available at BRIVIACT.com/tools-resources and learn how they can help make a difference for your child's treatment journey.



Seizure Diary: Allows you to record your child's seizures and provide a more complete picture for your doctor of where, when, and how often seizures occur



Caregiver Guide to Recording Seizures: Learn how you can help by recording the details of your child's seizures in order to assist the doctor in making a diagnosis and discussing treatment decisions



Other Epilepsy Resources: Find a list of organizations focused on epilepsy research, care, and support



SAVE ON BRIVIACT

Are you commercially insured? You could save on your child's BRIVIACT prescription.

A savings card works like a coupon—you can take it to your pharmacist and save money on your copay for your child's prescription.

Visit www.BRIVIACT.com to check eligibility and register for a savings card.



If you qualify,* you may pay as little as **\$10** for each 30-day supply of BRIVIACT.



*See BRIVIACT.com for additional Terms and Conditions.

HAVE MORE QUESTIONS?



Speak to a specialized nurse or pharmacist.

UCBCares® is a team of knowledgeable, experienced healthcare professionals (nurses and pharmacists) and customer service professionals who can guide you through questions you may have about BRIVIACT, savings, financial assistance, and more.

CONTACT

UCBCares®

About People

to talk to our team of nurses, pharmacists, and customer service professionals.



Call 1-844-599-CARE (2273)



Email ucbcares@ucb.com



or submit your question online at askucbcares.com



Interested in learning more about BRIVIACT?
Visit BRIVIACT.com



HELP GIVE THEM A CHILDHOOD FILLED WITH MORE MEMORIES AND FEWER PARTIAL-ONSET SEIZURES

INDICATION

BRIVIACT® (brivaracetam) CV is a prescription medicine used to treat partial-onset seizures in people 1 month of age and older.

IMPORTANT SAFETY INFORMATION

BRIVIACT may cause serious side effects including suicidal thoughts or actions; drowsiness; tiredness; dizziness; balance and coordination problems; mood and behavior changes including irritability, anxiety, and psychotic symptoms; and allergic reactions to ingredients in BRIVIACT.

The most common side effects of BRIVIACT include sleepiness, dizziness, feeling tired, and nausea and vomiting. Side effects of BRIVIACT in children 1 month to less than 16 years of age are similar to those seen in adults.

BRIVIACT is a federally controlled substance (CV) because it can be abused or lead to dependence.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to UCB, Inc. at UCBCares (1-844-599-CARE [2273]).

Please see Important Safety Information on pages 12-14 of this brochure and review the [Medication Guide on BRIVIACT.com](#).

BRIVIACT® and UCBCares® are registered trademarks of the UCB Group of Companies.
©2023 UCB, Inc., Smyrna, GA 30080. All rights reserved.
US-BR-2300097

Save on BRIVIACT. See inside for details.

BRIVIACT®
(brivaracetam)  